

## Medicinal cannabis: Rational guidelines for dosing

Gregory T Carter<sup>1\*</sup>, Patrick Weydt<sup>2</sup>, Muraco Kyashna-Tocha<sup>3</sup> & Donald I Abrams<sup>4</sup>

### Addresses

<sup>1</sup>Department of Rehabilitation Medicine

<sup>2</sup>Departments of Neurology and Laboratory Medicine

University of Washington School of Medicine

Seattle

WA 98531

USA

Email: gtcarter@u.washington.edu

<sup>3</sup>The Cyber Anthropology Institute

Seattle

WA 98145

USA

<sup>4</sup>San Francisco General Hospital

Division of Hematology/Oncology

Department of Medicine

University of California

San Francisco

CA 94110

USA

\*To whom correspondence should be addressed.

**IDrugs** 2004 **7**(5):

© The Thomson Corporation ISSN 1369-7056

*The medicinal value of cannabis (marijuana) is well documented in the medical literature. Cannabinoids, the active ingredients in cannabis, have many distinct pharmacological properties. These include analgesic, antiemetic, antioxidative, neuroprotective and anti-inflammatory actions, as well as modulation of glial cells and tumor growth regulation. Concurrent with all these advances in the understanding of physiological and pharmacological mechanisms of cannabis, there is a strong need for developing rational guidelines for dosing. This paper will review the known chemistry and pharmacology of cannabis and, on that basis, discuss rational guidelines for dosing.*

**Keywords** Cannabinoids, cannabis, dosing, marijuana, pharmacology

### Introduction and brief historical background

Possibly the first references to the medicinal use of cannabis are found in the Chinese pharmacopoeia of Emperor Shen-Nung, written in 2737 BC. This document recommended cannabis for analgesia, rheumatism, beriberi, malaria, gout and poor memory [1]. Eastern Indian documents in the Atharvaveda, dating to about 2000 BC, also refer to the medicinal use of cannabis [2]. Archeological evidence has been found in Israel indicating that cannabis was used therapeutically during childbirth as an analgesic [3]. This use of cannabis continued in the West until the mid-1880s and continues today in parts of Asia. In ancient Greece and Rome, both the Herbal of Dioscorides and the writings of Galen refer to the use of medicinal cannabis [4].

The medicinal use of cannabis in western medicine occurred much later. There is mention of it in a treatise by Culpepper written in medieval times. British East India Company surgeon William O'Shaughnessy introduced cannabis for medicinal purposes into the United Kingdom following his observations while working in India in the 1840s. He used it in a tincture for a wide range of uses, including analgesia [5], and Queen Victoria used cannabis for relief of dysmenorrhoea in the same era [6]. In 1937, against the advice of most of the medical community and much of the

American Medical Society, the federal government criminalized non-medical cannabis. Cannabis was removed from the United States Pharmacopoeia in 1942 but up until that time physicians were still able to write a prescription for it [7].

The physiological mechanisms and therapeutic value of cannabinoids continue to be well documented in the medical literature [6-36]. However, there has been little written on appropriate dosing regimens for the medicinal use of cannabis. With current and emerging laws allowing physicians in many areas of the world to recommend the use of cannabis to treat symptoms of certain diseases and medical conditions, there is a need for medical literature describing rational dosing guidelines. This paper will review the known chemistry and pharmacology of cannabis and then, on that basis, discuss rational guidelines for dosing.

## **Chemistry and pharmacology of cannabis**

Cannabis is a complex plant, with several existing phenotypes, each containing over 400 chemicals [14,15]; approximately 60 are chemically unique and classified as plant cannabinoids [11,15]. Naturally occurring cannabinoids are also produced in the human body [8]. The cannabinoids are 21 carbon terpenes, biosynthesized predominantly via a recently discovered deoxyxylulose phosphate pathway [16], and are lipophilic.  $\Delta^9$ -tetrahydrocannabinol (THC) and  $\Delta^8$ -THC appear to produce the majority of the psychoactive effects of cannabis.  $\Delta^9$ -THC, the active ingredient in dronabinol (Marinol) is the most abundant cannabinoid in the plant and this has led researchers to hypothesize that it is the main source of the drug's impact [15]. Dronabinol is available by prescription as a schedule III drug.

Other major plant cannabinoids include cannabidiol and cannabinol, both of which may modify the pharmacology of THC and have distinct effects of their own. Cannabidiol is the second most prevalent active ingredient in cannabis and may produce most of its effects at moderate, mid-range doses. Cannabidiol becomes THC as the plant matures and this THC over time breaks down into cannabinol. Up to 40% of the cannabis resin in some strains is cannabidiol [15]. The amount varies according to plant; some varieties of *Cannabis sativa* have been found to contain no cannabidiol [6]. As cannabidiol may help reduce anxiety symptoms, cannabis strains without cannabidiol may produce more panic or anxiogenic side effects. Cannabidiol may exaggerate some of the effects of THC, including increasing THC-induced euphoria, while attenuating others. Cannabidiol competitively slows THC metabolism in the liver. Consequently, a dose of THC combined with cannabidiol will create more psychoactive metabolites than the same dose of THC alone [14,15]. In mice, pretreatment with cannabidiol increased brain levels of THC by ~ 3-fold and there is strong evidence that cannabinoids can increase the brain concentration and pharmacological actions of other drugs [16,17]. Some researchers have proposed that many of the negative side effects of dronabinol, including sedation and altered mental activity, could be reduced by combining it with cannabidiol or possibly other non-psychoactive cannabinoids [8].

Cannabidiol breaks down to cannabinol as the plant matures [15]. Much less is known about cannabinol, although it appears to have pharmacological properties that are quite different from cannabidiol. Cannabinol has significant anticonvulsant, sedative and other pharmacological activities likely to interact with the effects of THC [14]. Cannabinol may induce sleep and may provide some protection against seizures for epileptics [15-17].

Two physiologically occurring lipids, anandamide (AEA) and 2-arachidonylglycerol (2-AG), have been identified as endogenous cannabinoids (endocannabinoids), although there are likely to be more [18]. The physiological roles of these endocannabinoids have been only partially clarified but available evidence suggests that they function as diffusible and short-lived intercellular messengers that modulate synaptic transmission. Recent studies have provided strong experimental evidence that endogenous cannabinoids mediate signals retrogradely from depolarized postsynaptic neurons to presynaptic terminals to suppress subsequent neurotransmitter release, driving the synapse into an altered state [18-20]. Signaling by the endocannabinoid system appears to represent a mechanism by which neurons can communicate backwards across synapses to modulate their inputs.

There are two known cannabinoid receptor subtypes. Subtype 1 (CB<sub>1</sub>) is expressed primarily in the brain whereas subtype 2 (CB<sub>2</sub>) is expressed primarily in the immune system [10,20]. Cannabinoid receptors constitute a major family of G protein-coupled, seven-helix transmembrane nucleotides, similar to the receptors of other neurotransmitters such as dopamine, serotonin and norepinephrine. In fact, they are the most abundant G protein-coupled receptor in the brain [8,10,11]. Activation of protein kinases may be responsible for some of the cellular responses elicited by the CB<sub>1</sub> receptor

[21].

Because of this biochemical complexity, characterizing the clinical pharmacology of cannabis is challenging. Further complicating the evaluation of cannabis is the variable potency of the plant material used in research studies. The concentration of THC and other cannabinoids in cannabis varies greatly depending on growing conditions, plant genetics and processing after harvest [19]. The highest concentrations of bioactive compounds are found in the resin exuded by the flowering female plants [18,19]. Leaf mixtures of cannabis have concentrations of THC ranging from 0.3 to 4% by weight [18-20]. However, cannabis today is typically distributed as flowers and can contain anything from 8 to  $\geq 25\%$  of THC. Thus, one gram of cannabis flowers would typically contain 80 to 250 mg of THC [19].

The clinical pharmacology of cannabis containing high concentrations of THC may differ from plant material containing small amounts of THC and higher amounts of the other cannabinoids. Moreover, the bioavailability and pharmacokinetics of inhaled cannabis are substantially different than when cannabis is ingested [17,18].

### **Clinical pharmacology**

Although it is a potent drug that may produce psychoactive effects, THC (and the other cannabinoids) have relatively low toxicity, and lethal doses in humans have not been described [23,24]. The theoretical LD<sub>50</sub> is estimated to be 1 to 20,000 or 1 to 40,000, using a single cannabis cigarette as a unit of dose. Conversely stated, a human would have to consume 20,000 to 40,000 times the amount of cannabis contained in one cigarette, in a short period of time, to achieve lethality. Using this as a basis, it has been estimated that it would require 1500 pounds of cannabis smoked in 15 min to induce a lethal effect [25].

Central effects of cannabinoids include disruption of psychomotor behavior, short-term memory impairment, intoxication, stimulation of appetite, antinociceptive actions (particularly against pain of neuropathic origin) and anti-emetic effects. Although there are signs of mild cognitive impairment in chronic cannabis users there is little evidence that such impairments are irreversible, or that they are accompanied by drug-induced neuropathology. A proportion of regular users of cannabis will develop some tolerance [37]. A study by Hart and colleagues demonstrated that acute cannabis smoking produced minimal effects on complex cognitive task performance in experienced cannabis users, while still subjectively providing a euphoric 'high' [38]. The potential medical applications of both natural and synthetic cannabinoids are currently being tested in a number of clinical trials.

### **Delivery system and pharmacokinetics**

Route of administration is an important determinant of the pharmacokinetics of the cannabinoids in cannabis, particularly absorption and metabolism [39-42]. Typically, cannabis is smoked as a cigarette weighing between 0.5 and 1.0 g. After combustion and inhalation, peak venous blood levels of 75 to 150 nanograms per milliliter (ng/ml) of plasma appear when smoking is finished [39,43,44]. The main advantage of smoking is rapid onset of effect and easy dose titration. When cannabis is smoked, cannabinoids in the form of an aerosol in the inhaled smoke are absorbed and delivered to the brain rapidly, as would be expected of a highly lipid-soluble drug [41,45].

A person's smoking behavior during an experiment is difficult for a researcher to control, and smoking behavior is not easily standardized, although some research protocols for standardization of smoking have been developed [44]. An experienced cannabis smoker can titrate and regulate dose to obtain the desired acute effects and to minimize undesired effects [46,47]. Each inhalation delivers a discrete dose of cannabinoids to the body. Inhalation volume changes with phase of smoking, tending to be highest at the beginning and lowest at the end of smoking a cigarette. Some studies found frequent users to have higher inhalation volumes than less frequent cannabis users. Heavy users could absorb as much as 27% of available THC, which may be twice as much as an infrequent user may absorb [47]. During smoking, as the cigarette length shortens, the concentration of THC in the remaining cannabis increases. Thus, each successive inhalation contains an increasing concentration of THC [47]. However, up to 40% of the available THC may be completely combusted in the process of smoking and may not be biologically available. Assays of cannabinoids in blood or urine after smoking can partially quantify dose actually absorbed, but the analytic procedures are methodologically demanding [47,48].

After smoking, venous blood levels of THC fall precipitously within minutes, and an hour later they are ~ 5 to 10% of the peak level [40,41,43,44]. Plasma clearance of THC is quite high at  $\geq 950$  ml/min which is essentially the rate of hepatic blood flow. However, the rapid disappearance of THC from blood is largely due to redistribution to other tissues in the body rather than cannabinoid metabolism [40,41]. Metabolism in most tissues is relatively slow. Slow release of cannabinoids from tissues and subsequent metabolism results in a long elimination half-time. The terminal half-life of THC is estimated to be from ~ 20 h to as long as 10 to 13 days, although reported estimates vary considerably and are likely to reflect the sensitivity of the measurement assay.

Smoking anything, including cannabis, is not healthful for the lungs and airway system [49,50]. A healthier option may be vaporization; because cannabinoids are volatile, they will vaporize at a temperature much lower than actual combustion [51]. Heated air can be drawn through cannabis and the active compounds will vaporize, and these can then be inhaled. Vaporization delivers the substance in a rapid manner that, like smoking, can be easily titrated to the desired effect [9]. Theoretically this removes most of the health hazards of smoking, although this has not yet been studied. Furthermore, there may be differing vaporization points for the individual cannabinoids. Vaporized cannabis may have differing concentrations and ratios of cannabinoids compared to smoked cannabis, although this also needs further study.

Cannabis can also be ingested orally or through a feeding tube. Orally ingested THC or cannabis has quite different pharmacokinetics than when it is inhaled. The onset of action is delayed and titration of dosing is more difficult [52-55]. Maximum THC and other cannabinoid blood levels are only reached 1 to 6 h after an oral dose, with a half-life of 20 to 30 h [52-55]. The same is true of dronabinol capsules, which contain only synthetic THC and none of the other cannabinoids [54]. When orally ingested, THC is broken down in the liver to the byproduct 11-hydroxy-THC, which also has potent psychoactive effects. This metabolite occurs at a much lower concentration when cannabis is inhaled. Thus, when THC (dronabinol or cannabis) is ingested orally, more sedation occurs because of the 11-hydroxy-THC psychoactive metabolite [54].

### **Metabolism, bioavailability and drug interactions**

Some inactive carboxy metabolites have terminal half-lives of 50 h to 6 days or more and thus serve as markers of prior cannabis use in urine tests [55,56]. Most of the absorbed THC dose is eliminated in feces, and ~ 33% is eliminated in urine. THC enters enterohepatic circulation and undergoes hydroxylation and oxidation to 11-nor-9-carboxy- $\Delta^9$ -THC (9-COOH-9-THC). The glucuronide is excreted as the major urine metabolite along with ~ 18 nonconjugated metabolites. Frequent and infrequent cannabis users are similar in the way that they metabolize THC [53].

THC bioavailability from smoked cannabis varies greatly among individuals and also depends on the composition of the specific cannabis preparation. Bioavailability can range from 1 to 27% with variable bioavailability resulting from significant loss of THC in side stream smoke, as well as variation in individual smoking behaviors. This includes incomplete absorption from inhaled smoke, metabolism in lung, and cannabinoid pyrolysis (ie, destruction by combustion).

Cannabinoids appear to partially inhibit the metabolism of drugs metabolized by the hepatic cytochrome P450 enzyme system [57-60]. Thus, the absorption or clearance of other drugs taken with cannabis may be slowed or hastened depending on timing and sequence of drug ingestion and past exposure. THC is highly bound to plasma proteins (97 to 99%) and is likely to interact with other highly bound drugs because of competition for binding sites on plasma proteins [61,62].

### **Dronabinol**

The Food and Drug Administration (FDA) first licensed and approved dronabinol in 1986 for the treatment of nausea and vomiting associated with chemotherapy. The indication was expanded in 1992 for the treatment of anorexia associated with weight loss in patients with AIDS wasting syndrome. In a randomized, double-blind, placebo-controlled, 6-week study involving 139 patients, dronabinol provided statistically significant improvement in appetite and non-statistically significant trends toward improved body weight and mood, and decreases in nausea [63]. In 1999, the United States Drug Enforcement Administration, in cooperation with the FDA, reclassified the scheduling status of dronabinol from a Schedule II (CII) to a Schedule III (CIII) controlled substance (for definitions of schedules refer to

<http://www.dea.gov/pubs/csa/812.htm>).

In 454 patients with cancer who received a total of 750 courses of treatment for various malignancies, dronabinol capsules provided complete or partial success in easing nausea and vomiting in 68% of patients given dronabinol (< 7 mg/m<sup>2</sup>/day) and 64% of patients given dronabinol (> 7 mg/m<sup>2</sup>/day) [64].

According to the manufacturer the prescribed dose of dronabinol for appetite stimulation is 2.5 mg twice-daily, to be taken before lunch and dinner. For nausea, vomiting, and pain the dosing is 5 mg/m<sup>2</sup>. If the 5 mg dose is ineffective, incremental increases of 2.5 mg, up to a maximum of 15 mg, is recommended. The same dose can be taken every 2 to 4 h for a maximum of four to six doses a day. Regardless of the clinical setting in which it is prescribed, the maximum total recommended dose of dronabinol is 15mg/m<sup>2</sup> four- to six-times daily or ~ 100 to 120 mg a day [65].

## Clinical trials

There are a limited number of well-performed clinical trials from which to draw succinct dosing regimens. Clinical trials have typically used cannabis cigarettes supplied by the NIDA (National Institute on Drug Abuse) containing 3.5 to 4.0% of THC by weight [59,66,67]. Recently, Abrams *et al* conducted an open-label study in patients with confirmed HIV neuropathy with persistent neuropathic pain [68]. All patients had prior experience of smoking marijuana but had ceased for 30 days prior to admission. After a 2-day lead-in period, patients smoked one cigarette containing 3.56% of THC three times/day for 7 days. A heat-capsaicin-induced experimental pain model was used to clarify the effects of the THC. Marijuana smoking caused a drop in pain score to 20/100 with ten of 16 patients experiencing a 30% reduction in average daily pain. An excellent correlation was noted in the response to the heat-capsaicin model, as 14 of 16 patients experienced a 30% reduction in the area of secondary hyperalgesia after smoking [68].

Wade *et al* compared plant-derived cannabis extracts to standard treatments for neurogenic symptoms unresponsive to standard treatment in a double-blind, randomized, placebo-controlled, cross-over trial with 2-week treatment periods [69]. There were 24 patients total, with diagnoses including multiple sclerosis (n = 18), spinal cord injury (n = 4), brachial plexus injury (n = 1), and limb amputation due to neurofibromatosis (n = 1). Whole-plant extracts of either THC only, cannabidiol only, a mixed cannabinoid extract of both THC and cannabidiol in a 1:1 ratio, or a matched placebo were self-administered by sublingual spray at doses determined by titration against symptom relief or unwanted effects within the range of 2.5 to 120 mg/24 h. The results demonstrated that pain relief associated with both THC and cannabidiol was significantly superior to placebo. The mixed cannabinoid extract, compared to placebo, was significantly superior in providing pain relief and improving bladder control, muscle spasms and spasticity. Side effects were rare. Three patients had transient hypotension and intoxication with rapid initial dosing of the THC extract.

## Deriving dosing recommendations and guidelines

Cannabis has many variables that do not fit well with the typical medical model for drug prescribing. If the plant is used, the variations are extreme. Plants vary immensely by phenotypes, and even the time of harvest affects which cannabinoids are present and in what percentages. One person may be much more sensitive than another, heavy smokers may get a different chemical 'smorgasbord' than light smokers and ingestion may alter bioavailability. The bulk of the research into cannabis has primarily examined THC, the other cannabinoids to a lesser degree, and the combinations hardly at all, although this is beginning to change. These combinations are very important to medicinal users of cannabis as a number of positive synergistic effects could be involved [70-72]. All of these points make it imperative that the dosing is highly individualized, so a patient-determined, self-titrated dosing model is recommended. This self-titration model is acceptable given the variables discussed above, as well as the low toxicity of cannabis. This construct is not unique to cannabis. There are other drugs that have relatively low toxicity and high dosing ceilings (gabapentin being one notable example), and are titrated to effect.

To facilitate an understanding of the determination of these guidelines, an estimate as to the actual amount of THC obtained by a patient when smoking different strengths of cannabis must be derived. As noted earlier, with smoking as the delivery, 40% of the active ingredients are lost in side stream or combustion, and a maximum of 27% of the remaining active ingredients can actually be absorbed by the patient. Given this, the maximum THC absorbed by a patient using 1 g of cannabis containing 10% of THC would be 16.3 mg.

The only form of cannabinoid that is available by a formal, dose-specific prescription is dronabinol. There are too many variables in the published clinical trials and case series with raw cannabis to use those studies as a basis for deriving doses. Therefore, we will use the dronabinol prescription guidelines as published by the manufacturer and accepted by the FDA as the basis for formulating our dosing recommendations for natural cannabis. It is critical to note that dronabinol is an oral preparation and contains only THC. Most medicinal cannabis patients use smoking as the route of delivery. As we have previously noted there are significant differences in pharmacokinetics between oral consumption and smoking. Furthermore, there are varying physiological effects when the other cannabinoid forms are present, as is the case with natural cannabis plant material. It is also not clear how the original dosing construct for dronabinol was arrived at, although we assume it was done through clinical testing for therapeutic benefit versus side effects. Despite these inherent limitations, these calculations do provide approximate dose equivalents by weight and are useful as long as one recognizes these limitations.

Applying the known pharmacokinetics of cannabis, as described above, to a conservative dronabinol dosing model of 2.5 to 60 mg/day, we calculated the following doses for cannabis containing these percentages of THC (Table 1).

**Table 1. Amount of cannabis equivalent to dronabinol (2.5 to 60 mg).**

% of THC in cannabis	Amount of cannabis required to obtain:			
	2.5 mg of THC	10 mg of THC	30 mg of THC	60 mg of THC
5% THC	0.60 g	1.24 g	3.70 g	7.40 g
10% THC	0.30 g	0.62 g	1.85 g	3.70 g
15% THC	0.16 g	0.41 g	1.23 g	2.46 g
20% THC	0.10 g	0.31 g	0.93 g	1.86 g
25% THC	0.08 g	0.25 g	0.75 g	1.50 g
30% THC	0.05 g	0.20 g	0.62 g	1.24 g

These derived figures lie closely within the range of reported amounts. In informal surveys from patients in Washington and California, the average reported consumption of cannabis by medicinal users typically ranges between 10 to 20 g of raw cannabis per week, or ~ 1.42 to 2.86 g/day of cannabis. The average strength of medical cannabis used by the patients who reported these doses was 15% THC. Thus these patients were actually absorbing between 34 and 68 mg/day of THC from the raw cannabis. The mean strength of medical cannabis in this study was ~ 19% THC, which corresponds to 44 to 88 mg/day of THC actually being consumed by the patient [72]. These figures are all within a similar range.

Our recommended doses are further reinforced by two of the studies that utilized smoked cannabis in a well-documented dosing regime. Chang *et al* studied the effects of smoked cannabis dosed at 10 mg/m<sup>2</sup> five-times-daily, equivalent to 87.5 mg of THC a day for an average-sized person. This would be the equivalent of 3.6g of cannabis containing 15% of THC [73]. Vinciguerra *et al* studied smoked cannabis dosed at 5 mg/m<sup>2</sup> four-times-daily, or 35 mg of THC a day for an average person. This is the equivalent of ~ 1.4g of cannabis containing 15% of THC [74]. For the purposes of these calculations, we assumed a 5'7" person weighing 140 lbs, with a body surface area of 1.75 meters squared.

These doses all fall within the medical cannabis guidelines allowed in the Canadian medical system. The Canadian medical allowance is for 1 to 12 g/day with an average of > 5 g. These doses are also very similar to the dosing range reported in a recent survey of patients who use cannabis to control symptoms of amyotrophic lateral sclerosis [75]. Thus, despite all of the noted variables, there is remarkable consistency among our derived doses and the reported doses from a number of different sources noted here.

A final comment should be made regarding physiological tolerance to cannabinoids. Tolerance does play a significant role in cannabis use and tolerance may develop to any of the various cannabinoids [76]. With regard to treating chronic, intractable pain, physicians will often prescribe increasingly larger doses of long-acting opioids as

patients develop tolerance. These patients are also generally given prescriptions of fast onset, short-acting opioids for 'breakthrough pain'. This is accepted practice, despite the fact that opioids, even in an opioid-dependent patient, still have the capacity to suppress breathing to the extent of inducing respiratory arrest. Long-term cannabis users can develop tolerance but, as previously discussed, there is essentially no risk for overdose. Thus, it is conceivable that a long-term cannabis user may require significantly larger amounts of cannabis to achieve a therapeutic effect. In addition, those who ingest cannabis may also require significantly higher amounts. Until more refined and purified cannabinoid preparations are available it will not be possible to derive a more specific or exact dosing schedule.

## Conclusions

We have outlined reasonable guidelines for dosing of medical cannabis, based on the known pharmacology. Our dosing model is primarily derived from dronabinol (THC), since that is the only clearly defined, FDA approved dosing paradigm currently available. However, our derived dosing schedule did match reasonably well with the amounts of natural cannabis reported by medical users. In using our dosing guidelines clinicians must be aware that THC is not the only clinically useful and pharmacologically active cannabinoid. The effects of THC are clearly modulated by other cannabinoids, which may have unique effects of their own. The clinician must also be aware of patient tolerance, and differing routes of intake and delivery systems, which can affect pharmacokinetics and bioavailability. Recognizing this, we recommend that our guidelines are used as a construct to allow the physician and patient to develop an individual, self-titration dosing paradigm. Given the current state of the known, published pharmacology of cannabis, this is the best dosing model that can be derived.

## Acknowledgments

Supported by Research and Training Center Grant HB133B980008 from the National Institute on Disability and Rehabilitation Research, Washington, DC, USA. The authors would like to acknowledge Dale Gieringer, Martin Martinez and Ethan Russo for their help in preparing this manuscript.

## References

- of outstanding interest
  - of special interest
1. Li HL: **An archaeological and historical account of cannabis in China.** *Econ Bot* (1974) **28**:437-448.
  2. Indian Hemp Drugs Commission: **Report of the Indian Hemp Drugs Commission, 1893-94.** Simla, India, Government Central Printing House (1894).
  3. Zias J, Stark H, Sellgman J, Levy R, Werker L, Breuer A, Mechoulam R: **Early medical use of cannabis.** *Nature* (1993) **363**(6426):215.
  4. Dioscorides, Pedanius. *The Greek herbal of Dioscorides.* Translated by Goodyer J, Gunther RWT. Hafner Publishing, London, UK (1968).
  5. O'Shaughnessy WB: **On the preparations of the Indian hemp, or gunjah (Cannabis indica): Their effects on the animal system in health, and their utility in the treatment of tetanus and other convulsive diseases.** *Trans Med Phys Soc Bengal* (1838-1840).
  6. Martinez M: **A Brief History of Marijuana.** In: *The New Prescription: Marijuana as Medicine.* Podrebarac F (Ed), Quick American Archives, Oakland, California, USA (2000):1-18.
    - *This well-referenced book provides a balanced view of historical, medical, and legal issues regarding medical marijuana.*
  7. Fankhauser M. **History of Cannabis in Western Medicine.** In: *Cannabis and Cannabinoids: Pharmacology,*

*Toxicology, and Therapeutic Potential*. Grotenhermen F, Russo E (Eds), The Haworth Integrative Healing Press, Oxford, UK (2002):37-49.

8. Carter GT, Weydt P: **Cannabis: Old medicine with new promise for neurological disorders**. *Curr Opin Invest Drugs* (2002) **3**(3):437-440.

9. Campbell VA: **Tetrahydrocannabinol-induced apoptosis of cultured cortical neurones is associated with cytochrome c release and caspase-3 activation**. *Neuropharmacology* (2001) **40**(5):702-709.

10. Valjent E, Pages C, Rogard M, Besson MJ, Maldonado R, Caboche J:  **$\Delta^9$ -tetrahydrocannabinol-induced MAPK/ERK and Elk-1 activation *in vivo* depends on dopaminergic transmission**. *Eur J Neurosci* (2001) **14**(2):342-352.

11. Friedman H, Klein TW, Newton C, Daaka Y: **Marijuana, receptors and immunomodulation**. *Adv Exp Med Biol* (1995) **373**:103-113.

12. Chen Y, Buck J: **Cannabinoids protect cells from oxidative cell death: A receptor-independent mechanism**. *J Pharmacol Exp Ther* (2000) **293**(3):807-812.

•• *An important study documenting the neuroprotective effects of cannabinoids.*

13. Guzman M, Sanchez C, Galve-Roperh I: **Control of the cell survival/death decision by cannabinoids**. *J Mol Med* (2001) **78**(11):613-625.

14. Akinshola BE, Chakrabarti A, Onaivi ES: ***In-vitro* and *in-vivo* action of cannabinoids**. *Neurochem Res* (1999) **24**(10):1233-1240.

15. Hampson AJ, Grimaldi M, Axelrod J, Wink D: **Cannabidiol and (-) $\Delta^9$ -tetrahydrocannabinol are neuroprotective antioxidants**. *Proc Natl Acad Sci USA* (1998) **95**(14):8268-8273.

•• *Another study confirming the neuroprotective effects of cannabinoids.*

16. Hampson AJ, Grimaldi M, Lolic M, Wink D, Rosenthal R, Axelrod J: **Neuroprotective antioxidants from marijuana**. *Ann NY Acad Sci* (2000) **899**:274-282.

•• *Similar to references 7 and 10, this study also confirms the neuroprotective effects of cannabinoids.*

17. Nagayama T, Sinor AD, Simon RP, Chen J, Graham SH, Jin K, Greenberg DA: **Cannabinoids and neuroprotection in global and focal cerebral ischemia and in neuronal cultures**. *J Neurosci* (1999) **19**(8):2987-2995.

•• *See comments for reference 16.*

18. Eshhar N, Striem S, Biegon A: **HU-211, a non-psychotropic cannabinoid, rescues cortical neurones from excitatory amino acid toxicity in culture**. *Neuroreport* (1993) **5**(3):237-240.

•• *See comments for reference 16.*

19. Hollister LE: **Marijuana and immunity**. *J Psychoact Drugs* (1988) **20**(1):3-8.

20. Carter GT, Rosen BS: **Marijuana in the management of amyotrophic lateral sclerosis**. *Am J Hospice Palliative Care* (2001) **18**(4):264-270.

21. Hansen HH, Schmid PC, Bittigau P, Lastres-Becker I, Berrendero F, Manzanares J, Ikonomidou C, Schmid HH, Fernandez-Ruiz JJ, Hansen HS: **Anandamide, but not 2-arachidonoylglycerol, accumulates during *in vivo* neurodegeneration**. *J Neurochem* (2001) **78**(6):1415-1427.



22. Panikashvili D, Simeonidou C, Ben-Shabat S, Hanus L, Breuer A, Mechoulam R, Shohami E: **An endogenous cannabinoid (2-AG) is neuroprotective after brain injury.** *Nature* (2001) **413**(6855):527-531.  
●● See comments for reference 16.
23. Ferraro L, Tomasini MC, Cassano T, Bebe BW, Siniscalchi A, O'Connor WT, Magee P, Tanganelli S, Cuomo V, Antonelli T: **Cannabinoid receptor agonist WIN 55,212-2 inhibits rat cortical dialysate  $\gamma$ -aminobutyric acid levels.** *J Neurosci Res* (2001) **66**(2):298-302.
24. Sinor AD, Irvin SM, Greenberg DA: **Endocannabinoids protect cerebral cortical neurons from *in vitro* ischemia in rats.** *Neurosci Lett* (2000) **278**(3):157-160.  
●● See comments for reference 16.
25. Young F: **In the Matter of Marijuana Rescheduling Petition.** US Department of Justice, DEA (1988):Docket No 86-22.
26. Gurley RJ, Aranow R, Katz M: **Medicinal marijuana: A comprehensive review.** *J Psychoact Drugs* (1998) **30**(2):137-147.
27. Robson P: **Therapeutic aspects of cannabis and cannabinoids.** *Br J Psychiatry* (2001) **178**:107-115.
28. Rice AS: **Cannabinoids and pain.** *Curr Opin Invest Drugs* (2001) **2**(3):399-414.  
● An excellent review article summarizing the analgesic effects of cannabinoids.
29. Di Marzo, Bisogno T, De Petrocellis L: **Endocannabinoids: New targets for drug development.** *Curr Pharm Des* (2000) **6**(13):1361-1380.
30. Tramer MR, Carroll D, Campbell FA, Reynolds DJ, Moore RA, McQuay HJ: **Cannabinoids for control of chemotherapy induced nausea and vomiting: Quantitative systematic review.** *Br Med J* (2001) **323**(7303):16-21.
31. Pertwee RG: **Cannabinoid receptor ligands: Clinical and neuropharmacological considerations, relevant to future drug discovery and development.** *Expert Opin Investig Drugs* (2000) **9**(7):1553-1571.  
●● An excellent review article outlining the future directions for drug development in cannabinoid-based medicines.
32. Renn E, Mandel S, Mandel E: **The medicinal uses of marijuana.** *Pharm Ther* (2000) **25**(10):536-524.
33. Hubbard JR, Franco SE, Onaivi ES: **Marijuana: Medical implications.** *Am Fam Phys* (1999) **60**(9):2583-2588.
34. Voth EA, Schwartz RH: **Medicinal applications of  $\Delta^9$ -tetrahydrocannabinol and marijuana.** *Ann Intern Med* (1997) **126**(10):791-798.
35. Baker D, Pryce G, Croxford JL, Brown P, Pertwee RG, Huffman JW, Layward L: **Cannabinoids control spasticity and tremor in a multiple sclerosis model.** *Nature* (2000) **404**(6773):84-87.
36. Consroe P, Musty R, Rein J, Tillery W, Pertwee R: **The perceived effects of smoked cannabis on patients with multiple sclerosis.** *Eur Neurol* (1997) **38**(1):44-48.
37. Abrams DI: **Medical marijuana: Tribulations and trials.** *J Psychoact Drugs* (1998) **30**(2):163-169.
38. Hart CL, van Gorp W, Haney M, Foltin RW, Fischman MW: **Effects of acute smoked marijuana on complex cognitive performance.** *Neuropsychopharmacology* (2001) **25**(5):757-765.

•• This study confirms the minimal side effects of cannabis use on cognitive performance, particularly in chronic users.

39. Adams IB, Martin BR: **Cannabis: Pharmacology and toxicology in animals and humans.** *Addiction* (1996) **91**(11):1585-1614.

40. Pertwee RG: **Sites and Mechanisms of Action.** In: *Cannabis and Cannabinoids: Pharmacology, Toxicology, and Therapeutic Potential.* Grotenhermen F, Russo E (Eds), The Haworth Integrative Healing Press, Oxford, UK (2002):73-81.

41. Agurell S, Halldin M, Lindgren JE, Ohlsson A, Widman M, Gillespie H, Hollister L: **Pharmacokinetics and metabolism of  $\Delta^1$ -tetrahydrocannabinol and other cannabinoids with emphasis on man.** *Pharmacol Rev* (1986) **38**(1):21-43.

42. Barnett G, Licko V, Thompson T: **Behavioral pharmacokinetics of marijuana.** *Psychopharmacology (Berl)* (1985) **85**(1):51-56.

43. Huestis MA, Henningfield JE, Cone EJ: **Blood cannabinoids. 1. Absorption of THC and formation of 11-OH-THC and THCCOOH during and after smoking marijuana.** *J Anal Toxicol* (1992) **16**(5):276-282.

44. Huestis MA, Sampson AH, Holicky BJ, Henningfield JE, Cone EJ: **Characterization of the absorption phase of marijuana smoking.** *Clin Pharmacol Ther* (1992) **52**(1):31-41.

45. Matthias P, Tashkin DP, Marques-Magallanes JA, Wilkins JN, Simmons MS: **Effects of varying marijuana potency on deposition of tar and  $\Delta^9$ -THC in the lung during smoking.** *Pharmacol Biochem Behav* (1997) **58**(4):1145-1150.

46. Heishman SJ, Stitzer ML, Yingling JE: **Effects of tetrahydrocannabinol content on marijuana smoking behavior, subjective reports, and performance.** *Pharmacol Biochem Behav* (1989) **34**(1):173-179.

47. Herning RI, Hooker WD, Jones RT: **Tetrahydrocannabinol content and differences in marijuana smoking behavior.** *Psychopharmacology* (1986) **90**(2):160-162.

48. Johansson E, Halldin MM, Agurell S, Hollister LE, Gillespie HK: **Terminal elimination plasma half-life of  $\Delta^1$ -tetrahydrocannabinol ( $\Delta^1$ -THC) in heavy users of marijuana.** *Eur J Clin Pharmacol* (1989) **37**(3):273-277.

49. Polen MR, Sidney S, Tekawa IS, Sadler M, Friedman GD: **Health care use by frequent marijuana smokers who do not smoke tobacco.** *West J Med* (1993) **158**(6):596-601.

50. Benowitz NL, Jones RT: **Cardiovascular and metabolic considerations in prolonged cannabinoid administration in man.** *J Clin Pharmacol* (1981) **21**(8-9 Suppl):214S-223S.

51. Gieringer D: **Cannabis vaporization: a promising strategy for smoke harm reduction.** *Journal of Cannabis Therapeutics* (2001) **1**(4):153-170.

52. Jones RT, Benowitz NL, Herning RI: **Clinical relevance of cannabis tolerance and dependence.** *J Clin Pharmacol* (1981) **21**(8-9 Suppl):143S-152S.

53. Kelly P, Jones RT: **Metabolism of tetrahydrocannabinol in frequent and infrequent marijuana users.** *J Anal Toxicol* (1992) **16**(4):228-235.

54. Grotenhermen F: **Effects of Cannabis and Cannabinoids.** In: *Cannabis and Cannabinoids: Pharmacology,*

*Toxicology, and Therapeutic Potential*. Grotenhermen F, Russo E (Eds), The Haworth Integrative Healing Press, Oxford, UK (2002):55-72.

55. Mechoulam R, Devane WA, Breuer A, Zahalka J: **A random walk through a cannabis field**. *Pharmacol Biochem Behav* (1991) **40**(3):461-464.

• *This paper delineates many of the complexities of the cannabis plant, including variations in species subtypes.*

56. Hollister LE: **Interactions of cannabis with other drugs in man**. In: *Strategies for Research on the Interactions of Drugs of Abuse*. National Institute on Drug Abuse Research Monograph 68. Braude MC, Ginzburg HM (Eds) DHHS Publication No (ADM)86-1453. Washington, DC, USA. Superintendent of Documents, US Government Printing Office (1986):110-116.

57. Fellermeier M, Eisenreich W, Bacher A, Zenk MH: **Biosynthesis of cannabinoids. Incorporation experiments with <sup>13</sup>C-labeled glucoses**. *Eur J Biochem* (2001) **268**(6):1596-1604.

58. Rueda D, Galve-Roperh I, Haro A, Guzman M: **The CB<sub>1</sub> cannabinoid receptor is coupled to the activation of c-Jun N-terminal kinase**. *Mol Pharmacol* (2000) **58**(4):814-820.

59. Kosel BW, Aweeka FT, Benowitz NL, Shade SB, Hilton JF, Lizak PS, Abrams DI: **The effects of cannabinoids on the pharmacokinetics of indinavir and nelfinavir**. *AIDS* (2002) **16**(4):543-550.

•• *This paper describes the hepatic metabolism of cannabinoids through the cytochrome P450 system.*

60. Benowitz NL, Jones RT: **Effect of  $\Delta^9$ -tetrahydrocannabinol on drug distribution and metabolism: Antipyrine, pentobarbital and ethanol**. *Clin Pharmacol Ther* (1977) **22**(3):259-268.

61. Gustafson RA, Levine B, Stout PR, Klette KL, George MP, Moolchan ET, Huestis MA: **Urinary cannabinoid detection times after controlled oral administration of delta9-tetrahydrocannabinol to humans**. *Clin Chem* (2003) **49**(7):1114-1124.

62. Benowitz NL, Nguyen T, Jones RT, Herning RI, Bachman J: **Metabolic and psychophysiologic studies of cannabidiol-hexobarbital interaction**. *Clin Pharmacol Ther* (1980) **28**(1):115-120.

63. Nelson K, Walsh D, Deeter P, Sheehan F: **A phase II study of  $\Delta^9$ -tetrahydrocannabinol for appetite stimulation in cancer-associated anorexia**. *J Palliat Care* (1994) **10**(1):14-18.

• *This is an important clinical trial defining the benefit of cannabis in appetite stimulation.*

64. Abrams DI, Hilton JF, Leiser RJ, Shade SB, Elbeik TA, Aweeka FT, Benowitz NL, Bredt BM, Kosel B, Aberg JA, Deeks SG *et al*: **Short-term effects of cannabinoids in patients with HIV-1 infection: A randomized, placebo-controlled clinical trial**. *Ann Intern Med* (2003) **139**(4):258-266.

•• *This is a major clinical trial looking at the effects of cannabinoids in the HIV-infected population.*

65. Unimed Pharmaceuticals: **Marinol Capsules product information package insert**. (2001).

66. Kelly TH, Foltin RW, Emurian CS, Fischman MW: **Effects of  $\Delta^9$ -THC on marijuana smoking, dose choice, and verbal report of drug liking**. *J Exp Anal Behav* (1994) **61**(2):203-211.

67. Ware MA, Doyle CR, Woods R, Lynch ME, Clark AJ: **Cannabis use for chronic non-cancer pain: Results of a prospective survey**. *Pain* (2003) **102**(1-2):211-216.

68. Jay C, Shade S, Vizoso H, Reda H, Petersen K, Rowbotham M, Abrams D: **The effect of smoked marijuana on chronic neuropathic and experimentally induced pain in HIV neuropathy: Results of an open-label pilot study**. *Int Conf Retroviruses Opportunistic Infect* (2004):Abs 496.

69. Wade DT, Robson P, House H, Makela P, Aram J: **A preliminary controlled study to determine whether whole-plant cannabis extracts can improve intractable neurogenic symptoms.** *Clin Rehabil* (2003) **17**(1):21-29.
70. Carlini EA, Karniol IG, Renault PF, Schuster CR: **Effects of marijuana in laboratory animals and man.** *Br J Pharmacol* (1974) **50**(2):299-309.
71. Fairbairn JW, Pickens JT: **Activity of cannabis in relation to its  $\Delta$ -trans-tetrahydro-cannabinol content.** *Br J Pharmacol* (1981) **72**(3):401-409.
72. Gieringer D: **Medical Cannabis Potency Testing Project.** *Bulletin of the Multidisciplinary Association for Psychedelic Studies* (1999) **9**(3):20-22.
73. Chang AE, Shiling DJ, Stillman RC, Goldberg NH, Seipp CA, Barofsky I, Simon RM, Rosenberg SA:  **$\Delta^9$ -tetrahydrocannabinol as an antiemetic in cancer patients receiving high-dose methotrexate. A prospective, randomised evaluation.** *Ann Intern Med* (1979) **91**(6):819-824.
74. Vinciguerra V, Moore T, Brennan E: **Inhalation marijuana as an antiemetic for cancer chemotherapy.** *New York State J Med* (1988) **88**(10):525-527.
75. Amtmann D, Weydt P, Johnson K, Jensen MP, Carter GT: **Survey of cannabis use in patients with amyotrophic lateral sclerosis.** *Am J Hosp Palliat Care* (2004) **21**(2):95-104.
76. Wilkinson JD, Whalley BJ, Baker D, Pryce G, Constanti A, Gibbons S, Williamson EM: **Medicinal cannabis: Is  $\Delta^9$ -tetrahydrocannabinol necessary for all its effects.** *J Pharm Pharmacol* (2003) **55**(12):1687-1694.